

# *Internet* Electronic Journal of **Molecular Design**

June 2005, Volume 4, Number 6, Pages 413–422

Editor: Ovidiu Ivanciuc

Proceedings of the Internet Electronic Conference of Molecular Design 2004  
IECMD 2004, November 29 – December 12, 2004

## **Numerical Evaluation of Two–Center Overlap Integrals Over Slater–Type Orbitals and Convergence Properties**

Hassan Safouhi and Ahmed Bouferguene

Faculté Saint–Jean/University of Alberta, 8406, 91street, Edmoton (AB), Canada T6C 4G9

Received: November 16, 2004; Revised: February 4, 2005; Accepted: March 23, 2005; Published: June 30, 2005

### **Citation of the article:**

H. Safouhi and A. Bouferguene, Numerical Evaluation of Two–Center Overlap Integrals Over Slater–Type Orbitals and Convergence Properties, *Internet Electron. J. Mol. Des.* **2005**, 4, 413–422, <http://www.biochempress.com>.

# Numerical Evaluation of Two–Center Overlap Integrals Over Slater–Type Orbitals and Convergence Properties<sup>#</sup>

Hassan Safouhi \* and Ahmed Bouferguene

Faculté Saint–Jean/University of Alberta, 8406, 91street, Edmoton (AB), Canada T6C 4G9

Received: November 16, 2004; Revised: February 4, 2005; Accepted: March 23, 2005; Published: June 30, 2005

*Internet Electron. J. Mol. Des.* 2005, 4 (6), 413–422

## Abstract

**Motivation.** Among the molecular integrals, the two–center overlap integrals play a major role in any accurate molecular structure calculation. They are central to the calculation of multicenter overlap integrals when using the series expansion formulae for Slater type functions about a new center. Consequently, these integrals require an accurate and fast numerical evaluation. Recently, we showed that these integrals are suitable to apply the nonlinear  $\bar{D}$  transformation of Sidi, which is shown to be highly efficient in improving convergence of highly oscillatory integrals.

**Method.** In this work, we present an algorithm for a numerical evaluation of the molecular integrals under consideration over STOs. Convergence properties in the numerical evaluation of these molecular integrals are discussed. It is now shown that the approximation obtained using the nonlinear  $\bar{D}$  transformation converges to the exact value of the integral without any constraint.

**Results.** Numerical results are obtained for two–center overlap integrals over Slater type orbitals with HCN, C<sub>2</sub>H<sub>2</sub>, BH<sub>3</sub> and CH<sub>4</sub> molecules. Comparisons with results obtained using the ACJU code developed by Homeier *et al.* are presented. Numerical results from the literature were also reproduced using the algorithm described in the present work.

**Conclusions.** The results obtained in this work illustrate the efficiency of the algorithm based on the nonlinear  $\bar{D}$  transformation, which will lead to a highly accurate algorithm for the numerical evaluation of the integrals under consideration.

**Keywords.** Molecular electronic integrals; Slater type orbitals; B functions; nonlinear transformations; convergence accelerators; numerical integration.

## Abbreviations and notations

STO, Slater–type orbital

ETO, exponential–type orbital

GTO, Gaussian–type orbital

## 1 INTRODUCTION

The numerical evaluation of two–center overlap integrals over exponential type functions is of great importance for any accurate molecular structure calculations. Multicenter molecular integrals can be expressed in terms of the two–center integrals, therefore the accurate and rapid numerical

<sup>#</sup> Presented in part at the Internet Electronic Conference of Molecular Design 2004, IECMD 2004.

\* Correspondence author; phone: 1–780–485–8631; fax: 1–780–465–8760; E–mail: hassan.safouhi@ualberta.ca.

evaluation of these integrals becomes more important in quantum mechanical calculations of the electronic structure of molecules.

STOs [1,2] are chosen for the expansion of atomic orbitals. These functions have a dominating position among ETOs, due to the fact that their analytical expression is very simple. STOs are better suited than GTOs to represent electron wave functions near the nucleus and at long range, provided that multicenter integrals using such functions could be computed efficiently.

STOs can be expressed as finite linear combinations of B functions [3,4,5]. The Fourier transforms of these B functions are exceptionally simple [6,7] and well adapted to the Fourier transform method [8,9], which led to analytical expressions for multicenter electronic integrals over B functions. These analytical expressions involve two-dimensional integral representations, which present severe numerical and computation difficulties. The integrand of the inner semi-infinite integral is a very oscillating function due to the presence of spherical Bessel function.

The molecular integrals under consideration are to be evaluated via a numerical quadrature of integral representations in terms of nonphysical integration variables. These integral representations were derived with the help of the Fourier transformation Method.

The semi-infinite integrals can be transformed into infinite series of integrals of alternating sign. These series are slowly convergent and this is why their use is prohibitively long for sufficient accuracy. The epsilon algorithm of Wynn [10] or Levin's  $u$  transform [11], accelerate the convergence of infinite series but in the case of the semi-infinite integrals involved in the analytical expressions of molecular integrals, the calculation times for a sufficient accuracy still long. Therefore new numerical integration techniques are required.

Recently [12], we demonstrate that the semi-infinite integrals under consideration are suitable to apply the nonlinear  $\bar{D}$  transformation of Sidi [13,14,15]. It is shown that the  $\bar{D}$  transformation is much more efficient and rapid compared with the alternative cited above.

In the present work, we discussed the convergence properties of  $\bar{D}$  transformation in improving convergence of the semi-infinite integrals occurring in the analytic expression of overlap integrals and we also presented the algorithm for the numerical evaluation of overlap integrals over STOs. We performed calculations with HCN, C<sub>2</sub>H<sub>2</sub>, BH<sub>3</sub> and CH<sub>4</sub> molecules and we reproduce values from table I in [16]. We also used the ACJU code developed Homeier *et al.* [17] to perform the same calculations. From the numerical tables, one can easily notice that our numerical results are in a complete agreement with those listed in Table I in [16] and with those obtained using the ACJU code. This illustrates clearly that the approach using nonlinear transformations for improving convergence of oscillatory integrals will probably lead to an efficient package for accurate numerical evaluation of all molecular integrals over exponential type functions.

## 2 DEFINITIONS AND PROPERTIES

Slater-type orbital (STOs) are given by [1,2]:

$$\chi_{n,l}^m(\zeta, r) = \sqrt{\frac{(2\zeta)^{2n+1}}{(2n)!}} r^{n-1} e^{-\zeta r} Y_L^M(\theta_r, \varphi_r) \quad (1)$$

where  $Y_L^M(\theta_r, \varphi_r)$  stands for the spherical harmonic.

The B function is defined as follows [4,5]:

$$B_{n,l}^m(\zeta, r) = \frac{(\zeta r)^l}{2^{n+1}(n+1)!} \hat{k}_{n+\frac{1}{2}}(\zeta r) Y_L^M(\theta_r, \varphi_r) \quad (2)$$

where  $\hat{k}_{n+\frac{1}{2}}(\zeta r)$  stands for the reduced Bessel function.

The two-center overlap integral over STOs is defined by:

$$S = \left\langle \chi_{n_1,l_1}^{m_1}(\zeta_1, r_1) \middle| \chi_{n_2,l_2}^{m_2}(\zeta_2, r - R) \right\rangle \quad (3)$$

Using the fact that STFs can be expressed in terms of B functions, one can express the above integrals in terms of integrals over B functions, which are given by:

$$\hat{S} = \left\langle B_{n_1,l_1}^{m_1}(\zeta_1, r_1) \middle| B_{n_2,l_2}^{m_2}(\zeta_2, r - R) \right\rangle \quad (4)$$

The Fourier transform method allowed analytic expressions to be developed for the above integrals over B functions [18]. These analytic expressions involve semi-infinite highly oscillatory, which are the principal source of the difficulty in the numerical evaluation of the molecular integrals under consideration. These semi-infinite integrals are given by [18]:

$$I = \int_0^\infty \frac{x^{n_x}}{(\zeta_1^2 + x^2)^{k_1} (\zeta_2^2 + x^2)^{k_2}} j_\lambda(Rx) dx \quad (5)$$

Where  $n_x$ ,  $\lambda$ ,  $k_1$  and  $k_2$  are positive integers and they depend on the quantum numbers.  $j_\lambda(Rx)$  stands for the spherical Bessel function. In practice  $n_x, k_1, k_2, \lambda \leq 10$ .

Recently [12], we demonstrate the applicability of the nonlinear  $\bar{D}$  transformation of Sidi [13,14]. Numerical results were obtained for two-center overlap integrals over B functions and they were in a complete agreement with results in the literature [16] and with results obtained using existing codes such as ACJU developed by Homeier *et al.* [17].

In the present work, we demonstrate by using previous work of Sidi [13,19,20], that the approximation of the above semi-infinite integrals obtained using the nonlinear  $\bar{D}$  transformation converges to the exact value of the semi-infinite integrals without any constraint.

Calculations of these integrals over STFs were also performed and presented, to show that our approach will definitely lead to a highly efficient algorithm for the numerical evaluation of two-center overlap integrals over B functions and over STFs.

### 3 NONLINEAR TRANSFORMATION AND CONVERGENCE PROPERTIES

The approximation of the semi-infinite integral (5) using  $\bar{D}$  is given by:

$$\bar{D}_n^{(2)} = \int_0^{x_l} F(t) dt + x_l^2 g(x_l) j'_\lambda(Rx_l) \sum_{i=0}^{n-1} \frac{\bar{\beta}_{1,i}}{x_l^i} \quad (6)$$

where  $l=0,1,\dots,n$  and  $x_l$  are the successive positive zeros of the spherical Bessel function.  $F(t)$  is the integrand of the semi-infinite integral (5) and  $g(x)$  is given by:

$$g(x) = \frac{x^{n_x}}{(\zeta_1^2 + x^2)^{k_1} (\zeta_2^2 + x^2)^{k_2}} \quad (7)$$

The unknowns  $\bar{\beta}_{1,i}$  for  $i=0,1,\dots,n-1$  stand for the approximations of the coefficients of the asymptotic expansion of the semi-infinite integral (5) [23].

Let  $\Psi(x)$  be defined by:

$$\Psi(x) = x^2 g(x) j'_\lambda(Rx) \quad (8)$$

Let  $(\gamma_0, \gamma_1, \dots, \gamma_n)$  the first row of the inverse matrix of the linear system given by Eq. (6). From the fact that the first column of the matrix of the system (6) is the vector  $(1,1,\dots,1)^T$  it follows that:

$$\sum_{i=0}^n \gamma_i = 1 \quad (10)$$

In [13], Sidi demonstrated that if:

$$\sum_{i=0}^n |\gamma_i| \leq L < \infty \quad (11)$$

then the approximation  $\bar{D}_n^{(2)}$  converges to the exact value of the semi-infinite integral without any constraint:

$$\left| I - \bar{D}_n^{(2)} \right| = o(n^{-j}) \quad (12)$$

for any  $j > 0$ .

In [19], Sidi demonstrated that a necessary and sufficient for  $\gamma_i > 0, i = 0,1,\dots,n$ , is:

$$\Psi(x_i) \Psi(x_{i+1}) < 0 \quad (13)$$

From the fact that  $x_i$  are the leading positive zeros of the spherical Bessel function, it follows that:

$$j'_\lambda(Rx_i) j'_\lambda(Rx_{i+1}) < 0 \quad (14)$$

Now by using the fact that  $x^2 g(x) > 0$ , it follows that (13) is satisfied and consequently, the approximation  $\bar{D}_n^{(2)}$  converges to the exact value of the semi-infinite integral without any constraint.

#### 4 RESULTS AND DISCUSSION

The computation of the approximation  $\bar{D}_n^{(2)}$  (6), require the computation of the first derivative of the spherical Bessel function. This can be obtained using the following equations:

$$\frac{\lambda}{2\lambda + 1} j_{\lambda-1}(x) - \frac{\lambda - 1}{2\lambda + 1} j_{\lambda+1}(x) = j'_\lambda(x) \quad (15)$$

$$j_{\lambda-1}(x) - \frac{\lambda + 1}{x} j_\lambda(x) = j'_\lambda(x) \quad (16)$$

In our algorithm, we used the following procedure. Note that if  $x$  is a zero of the spherical Bessel function of order  $\lambda$  for  $\lambda \geq 1$ , then Eq. (16) becomes:

$$j_{\lambda-1}(x) = j'_\lambda(x) \quad (17)$$

From the above equation, it follows that it is faster to use Eq. (17) than Eq. (15). But in this case, one has to separate the case where  $\lambda = 1$ , where the use of Cramer's rule is possible since the zeros of  $j_0(x) = \frac{\sin(x)}{x}$  are equidistant [13]. In this case, the approximation is given by:

$$\bar{D}_n^{(2)} = \frac{\sum_{i=0}^n C_n^i (1+i)^n \left( \int_0^{x_i} F(t) dt \right) / [x_i^2 g(x_i)]}{\sum_{i=0}^n C_n^i (1+i)^n / [x_i^2 g(x_i)]} \quad (18)$$

where  $x_i$  are the successive positive zeros of the sine function and where  $C_n^i = \frac{n!}{i!(n-i)!}$ .

We evaluate two-center overlap integrals over STOs. First, we expressed these integrals over STOs in terms of integrals over B functions, by expressing STOs as finite linear combinations of B functions [21]. Then we used the analytic expression obtained by Weniger *et al.* for the two-center overlap integrals over B functions. These analytic expressions involve the semi-infinite integrals

given by Eq. (5). We notice that these semi-infinite integrals have compact analytic expressions in the case where the scaling parameters are equal [18]. In the case where the scaling parameters are not equal, the semi-infinite integrals are evaluated using the nonlinear  $\bar{D}$  transformation (6).

**Table 1.** Exponents of STOs for a series of orbitals

Orbitals	B	C	N	H
1s	4.649767	5.636105	6.621925	1.00000
2s	1.076139	1.346562	1.612481	
2p	1.226030	1.581274	1.929475	

**Table 2.** The geometry used for the molecular calculations

Molecules	Geometry	Cartesian Coordinates
C <sub>2</sub> H <sub>2</sub>	Linear	H <sup>1</sup> (0.0, 0.0, -a-b/2)
	H-C = a = 2.002 a.u.	C <sup>1</sup> (0.0, 0.0, -b/2)
	C-C = b = 2.281 a.u.	C <sup>2</sup> (0.0, 0.0, b/2)
		H <sup>2</sup> (0.0, 0.0, a+b/2)
BH <sub>3</sub>	Equilateral Triangle, Planar	B ( 0.0, 0.0, 0.0)
	B-H = a = 2.250 a.u.	H <sup>1</sup> ( 0.0, 0.0, a)
	b = 3.897 a.u.	H <sup>2</sup> ( b/2, 0.0, -a/2)
		H <sup>3</sup> (-b/2, 0.0, -a/2)
CH <sub>4</sub>	Regular Tetrahedron	C ( 0.0, 0.0, 0.0)
	C-H = a = 2.0665 a.u.	H <sup>1</sup> ( b, b, b)
	b = 1.1931 a.u.	H <sup>2</sup> (b,-b, -b)
		H <sup>3</sup> (-b,b, -b)
		H <sup>4</sup> (-b,-b, b)
HCN	Linear	H(0.0, 0.0, -a)
	H-C = a = 2.000 a.u.	C(0.0, 0.0, 0.0)
	C-N = b = 2.187 a.u.	N(0.0, 0.0, b)

**Table 3.** Values obtained for the two-center overlap integrals over STOs

Molecules	Integrales	Values <sup><math>\bar{D}</math></sup>	Values <sup>ACJU</sup>
C <sub>2</sub> H <sub>2</sub>	$\langle 1s^{C1}   1s^{C2} \rangle$	0.17999067797259(-3)	0.179990672868803(-3)
	$\langle 2s^{C1}   1s^{C2} \rangle$	0.75773470045080(-1)	0.757734700451693(-1)
	$\langle 2p_z^{C1}   1s^{C2} \rangle$	0.11347210396549( 0)	0.113472103965470( 0)
	$\langle 2p_z^{C1}   2p_z^{C2} \rangle$	-0.28034732991369( 0)	-0.280347329913672( 0)
	$\langle 2p_{+1}^{C1}   2p_z^{C2} \rangle$	0.35111604505883( 0)	0.351116045058850( 0)
BH <sub>3</sub>	$\langle 1s^B   1s^{H1} \rangle$	0.84488384216158(-1)	0.844883842160902(-1)
	$\langle 2s^B   1s^{H1} \rangle$	0.61417323135024( 0)	0.614173231350241( 0)
	$\langle 2p_z^B   1s^{H1} \rangle$	0.54664981962486( 0)	0.546649819624870( 0)
	$\langle 1s^{H1}   1s^{H2} \rangle$	0.20218404614891( 0)	0.202184046148936( 0)
CH <sub>4</sub>	$\langle 1s^C   1s^{H1} \rangle$	0.75634817563177(-1)	0.756348175631746(-1)
	$\langle 2s^C   1s^{H1} \rangle$	0.62274411537583( 0)	0.622744115375811( 0)
	$\langle 2p_z^C   1s^{H1} \rangle$	0.47623357260361( 0)	0.274991804720407( 0)
	$\langle 1s^{H1}   1s^{H2} \rangle$	0.27969702358746( 0)	0.279697023587438( 0)
HCN	$\langle 1s^N   1s^C \rangle$	0.12627258447390(-3)	0.126272578207178(-3)
	$\langle 2s^N   1s^C \rangle$	0.73192433120423(-1)	0.731924331205254(-1)
	$\langle 2p_z^N   1s^C \rangle$	-0.99048113986006(-1)	-0.991971925002319(-1)

For the computation of the function  $F(x)$ , we transform the finite integral as follows:

$$\int_0^{x_n} F(t) dt = \sum_{l=0}^{n-1} \int_{x_l}^{x_{l+1}} F(t) dt \quad (19)$$

For the numerical evaluation of each term of the finite sum in the right hand side of Eq. (19), we used Gauss–Legendre quadrature of order 48. The linear system given by Eq. (6) is solved using the LU decomposition method.

**Table 4.** Values obtained for two–center overlap integrals over STOs. The two–centers are separated by unit distance in the z direction. Results are computed in double precision. Numbers in parenthesis represent powers of 10.

$n_1$	$l_1$	$m_1$	$\xi_1$	$n_2$	$l_2$	$m_2$	$\xi_2$	Values $\bar{D}$	Values ACJU	Values [16]
1	0	0	0.1	1	0	0	0.1	0.998337272657	0.998337285	0.998337285
1	0	0	1.0	1	0	0	0.1	0.187051615915	0.187051616	0.187051616
1	0	0	5.0	1	0	0	0.1	0.201680371415(-1)	0.20168086523(-1)	0.2016803717(-1)
1	0	0	1.0	1	0	0	1.0	0.858385362733	0.858385363	0.858385363
1	0	0	5.0	1	0	0	1.0	0.239940017735	0.239940018	0.239940018
1	0	0	5.0	1	0	0	5.0	0.965772403202(-1)	0.96577240320(-1)	0.9657724032(-1)
8	0	0	1.0	8	0	0	1.0	0.989015721	0.989015721	0.989015721
8	0	0	5.0	8	0	0	1.0	0.107437341700(-1)	0.10743703415(-1)	0.1074373417(-1)
8	0	0	5.0	8	0	0	5.0	0.78523085	0.78523085	0.78523085
5	4	0	1.0	5	4	0	0.1	0.219794514202(-2)	0.22132767718(-2)	0.2213276772(-2)
5	4	4	1.0	5	4	4	0.1	0.225942860190(-2)	0.22596477308(-2)	0.2259647731(-2)
5	4	0	5.0	5	4	0	0.1	0.664469344963(-6)	0.66675704065(-6)	0.6667585746(-6)
5	4	4	5.0	5	4	4	0.1	0.729423185992(-6)	0.72945439326(-6)	0.7294559035(-6)
5	4	0	1.0	5	4	0	1.0	0.768617011	0.768617016	0.768617016
5	4	4	1.0	5	4	4	1.0	0.955778746	0.955778746	0.955778746
5	4	0	5.0	5	4	0	1.0	0.900262308903(-2)	0.90026230924(-2)	0.9002623092(-2)
5	4	4	5.0	5	4	4	1.0	0.318003745747(-1)	0.31800374574(-1)	0.3180037457(-1)
5	4	0	5.0	5	4	0	5.0	-0.138257012	-0.138257012	-0.138257012
5	4	4	5.0	5	4	4	5.0	0.356825987	0.356825987	0.356825987
1	0	0	0.1	1	0	0	0.1	0.998337273	0.998337285	0.998337285

For the numerical evaluation of Gaunt coefficients, which occur in the analytic expression of the molecular integrals, we used the subroutine GAUNT.F developed by Weniger [22]. The spherical harmonics  $Y_l^m(\theta_r, \varphi_r)$  are computed using the recurrence formulae presented in [22].

We used the ACJU program developed by Homeier *et al.* [17] to evaluate the molecular integrals under consideration. As it can be seen from the numerical tables our numerical results are in a complete agreement with those obtained using ACJU program.

Table 1 contains the values of the screening parameters, which occur in the analytic expression of STOs. Table 2 contains the geometry used for the calculations that we performed for the present work. Table 3 contains values obtained for two–center overlap integrals over STOs with HCN,

C<sub>2</sub>H<sub>2</sub>, BH<sub>3</sub> and CH<sub>4</sub> molecules. The ACJU code was also used to perform the same calculations. Table 4 contains values obtained for two-center overlap integrals over STOs. In this table, we listed the results obtained using the ACJU code and the results obtained by Talman [16].

As it can be seen from this table our results are in a complete agreement with those obtained using the ACJU code and with the results obtained by Talman [16]. The abbreviations 2p<sub>z</sub> and 2p<sub>+1</sub> refer to orbitals defined with the quantum numbers:  $n = 2, l = 1, m = 0$  and  $n = 2, l = 1, m = 1$  respectively. Numbers in parentheses represent powers of 10. All the calculations were performed on a PC-Workstation Intel Xeon Processor 2.4GHz.

## 5 CONCLUSIONS

Analytic expressions for the two-center overlap integrals over STOs are obtained by expressing these STOs in terms of the so-called B functions and then by applying the Fourier transform method. These analytic expressions turned out very difficult to evaluate to because of the presence of highly semi-infinite integrals involving spherical Bessel functions.

It was shown that these semi-infinite integrals are suitable to apply the nonlinear  $\bar{D}$  method, which consists on transforming the semi-infinite integrals involving Bessel functions into asymptotic expansions in inverse powers of  $x$  as  $x \rightarrow \infty$ . These asymptotic expansions are transformed into sets of linear equations. The approximations of semi-infinite integrals are obtained by solving these linear systems and it is shown that the approximations obtained using  $\bar{D}$  transformation converge to the exact values of the semi-infinite integrals.

The numerical results show that the approach described in this work yields values for these integrals to a high pre-determined accuracy and in a complete agreement with values from the literature. These results confirm that this  $\bar{D}$  transformation represents another most significant advance in molecular multicenter integral problems.

## Acknowledgment

The first author acknowledges the financial support of this research by the Natural Sciences and Engineering Research Council of Canada (NSERC).

## 6 REFERENCES

- [1] J. C. Slater, Atomic shielding constants, *Phys. Rev.* **1930**, 36, 57.
- [2] J. C. Slater, Analytic atomic wave functions, *Phys. Rev.* **1932**, 42, 33.
- [3] Shavitt, The Gaussian function in calculation of statistical mechanics and quantum mechanics, *Methods in Computational Physics.2. Quantum.Mechanics*, edited by B. Alder, S. Fernbach, M. Rotenberg, (Academic Press,

- New York), 1963.
- [4] Filter and E. O. Steinborn, Extremely compact formulas for molecular one–electron integrals and Coulomb integrals over Slater–type orbitals, *Phys. Rev. A* **1978**, 18, 1.
- [5] E. O. Steinborn and E. Filter, Translations of fields represented by spherical–harmonics expansions for molecular calculations. III. Translations of reduced Bessel functions, Slater–type s–orbitals, and other functions, *Theor. Chim. Acta.* **1975**, 38, 273.
- [6] J. Weniger, Reduzierte Bessel–Funktionen als LCAO–Basissatz: Analytische und numerische Untersuchungen, Ph.D. thesis, Universität Regensburg, 1982.
- [7] E. J. Weniger and E. O. Steinborn, The Fourier transforms of some exponential–type functions and their relevance to multicenter problems, *J. Chem. Phys.* **1983**, 78, 6121.
- [8] P. Trivedi and E. O. Steinborn, Fourier transform of a two–center product of exponential–type orbitals. Application to one– and two–electron multicenter integrals, *Phys. Rev. A.* **1983**, 27, 670.
- [9] J. Grotendorst and E. O. Steinborn, Numerical evaluation of molecular one– and two–electron multicenter integrals with exponential–type orbitals via the Fourier–transform method, *Phys. Rev. A.* **1988**, 38, 3857.
- [10] D. Levin, Developement of non–linear transformations for improving convergence of sequences, *Int. J. Comput. Math.* **1973**, B 3, 371.
- [11] P. Wynn, On a device for computing the  $e_m(S_n)$  transformation, *Math. Tables Aids Comput.* **1956**, 10, 91.
- [12] H. Safouhi and L. Berlu, Nonlinear transformations for accurate numerical evaluation of two–center overlap and Coulomb integrals over B functions, *J. Phys. A: Math. Gen.* **2004**, submitted.
- [13] A. Sidi, Extrapolation methods for oscillatory infinite integrals, *Inst. Maths. Applics.* **1980**, 26, 1.
- [14] A. Sidi, Computation of infinite integrals involving Bessel functions of arbitrary order by the  $\bar{D}$ –transformation, *J. Comp. Appl. Math.*, **1997**, 78, 125.
- [15] A. Sidi, Computation of infinite integrals involving Bessel functions of arbitrary order by the  $\bar{D}$ –transformation, *J. Comp. Appl. Math.* **1997**, 78, 125.
- [16] J. D. Talman, Expression for overlap integrals of Slater orbitals, *Phys. Rev. A.* **1993**, 48, 243.
- [17] H. H. H. Homeier, E. J. Weniger and E. O. Steinborn, Programs for the evaluation of overlap integrals with B functions, *Comput. Phys. Commun.* **1992**, 72, 269.
- [18] E. J. Weniger and E. O. Steinborn, The Fourier transforms of some exponential–type functions and their relevance to multicenter problems, *J. Chem. Phys.* **1983**, 78, 6121.
- [19] A. Sidi, An algorithm for a special case of a generalization of the Richardson extrapolation process, *Numer. Math.* **1982**, 299.
- [20] A. Sidi, Some properties of a generalization of the Richardson extrapolation process, *J. Inst. Maths. Applics.* **1979**, 327.
- [21] E. Filter and E. O. Steinborn, Extremely compact formulas for molecular one–electron integrals and Coulomb integrals over Slater–type orbitals, *Phys. Rev. A.* **1978**, 18, 1.
- [22] E. J. Weniger and E. O. Steinborn. Programs for the coupling of spherical harmonics, *Comput. Phys. Commun.* **1982**, 25, 149.
- [23] D. Levin and A. Sidi, Two new classes of non–linear transformations for accelerating the convergence of infinite integrals and series, *Appl. Math. Comput.* 1981, 9.

## Biographies

**Hassan Safouhi** is associate professor of mathematics at the Faculté Saint–Jean/University of Alberta. After obtaining a Ph.D. degree in applied mathematics from the University Blaise Pascal, Dr. Safouhi undertook postdoctoral research with Prof. André Joyal at the University of Québec at Montréal. More recently, Dr. Safouhi has collaborated on projects with the group “Theoretical Chemistry” of the University Blaise Pascal. Dr. Safouhi is dedicated to the development of new mathematical techniques for a rapid and accurate numerical evaluation of molecular multicenter integrals over Slater–type orbitals and related functions.

**Ahmed Bouferguene** is assistant professor of mathematics at Faculté Saint Jean, Univ. of Alberta. After a PhD devoted mainly to study numerical algorithms geared towards an efficient evaluation of molecular integrals over Slater Type Orbitals (STOs), he developed a software package based on STOs for molecular structure determination. He then

spent several years at Florida A&M University as a postdoctoral fellow during which he collaborated with Pr. H. W. Jones and Pr. C. A. Weatherford on various problems including multicenter integrals over STOs, electron/molecule scattering and molecules subjected to strong magnetic fields. At Ottawa University, the author collaborated with Pr. D. M. Bishop on a non-linear optics related problem the result of which was a technique allowing one to visualize the overall molecular polarizability and hyper-polarizability as a “sum” of “elementary” contributions from the functional groups of the molecule.